


1/21


FIGURE 1



Typical log-normal probability plot of particle size distribution for M9 preparation with total polymer concentration of 8% in the organic solvent.

2/21


FIGURE 2



Theophylline release profiles in HCl solution, pH 1.2 (first 2 hours) phosphate buffer, pH 7.5 (after 2 hours) at 37°C for M3 microspheres.

3/21


FIGURE 3



Theophylline release profiles in HCl solution, pH 1.2 (first 2 hours) and phosphate buffer, pH 7.5 (after 2 hours) at 37°C for M5 microspheres.

4/21


FIGURE 4



Theophylline release profiles in HCl solution, pH 1.2 (first 2 hours) and phosphate buffer, pH 7.5 (after 2 hours) at 37°C for M6 microspheres.

5/21


FIGURE 5



Theophylline release profiles in HCl solution, pH 1.2 (first 2 hours) and phosphate buffer, pH 7.5 (after 2 hours) at 37°C for M8 microspheres.

6/21


FIGURE 6



Theophylline release profiles in HCl solution, pH 1.2 (first 2 hours) and phosphate buffer, pH 7.5 (after 2 hours) at 37°C for M9 microspheres.

7/21


FIGURE 7



Theophylline release profiles in HCl solution, pH 1.2 (first 2 hours) and phosphate buffer, pH 7.5 (after 2 hours) at 37°C for M10 microspheres.

8/21


FIGURE 8



Theophylline release profiles from 180-250 μm size fraction microspheres for different microsphere preparations in HCl solution, pH 1.2 (first 2 hours) and phosphate buffer, pH 7.5 (after 2 hours) at 37°C.

9/21


FIGURE 9



Theophylline release profiles from 250-355 μm size fraction microspheres for different microsphere preparations in HCl solution (pH 1.2 first 2 hours) and phosphate buffer (pH 7.5) at 37°C.

10/21


FIGURE 10



Theophylline release profiles from 355-500 μm size fraction microspheres for different microspheres preparations in HCl solution (pH 1.2 first 2 hours) and phosphate buffer (pH 7.5) at 37°C.

11/21


FIGURE 11



Higuchi plots for dissolution of different size fractions for M9 microspheres phosphate buffer (pH 7.5 after 2 hours in HCl) at 37°C.

12/21


FIGURE 12



Theophylline release profiles in phosphate buffer pH 7.5 at 37°C for M3 microsphere preparation.

13/21


FIGURE 13



Typical log-probability plot of particle size distribution for K7 (5% HPC) preparation with total polymer concentration of 7.5% in the organic solvent.

14/21


FIGURE 14



Release profile of theophylline in simulated intestinal fluid from K3 (1% HPC) microspheres with a total polymer concentration of 7.5% in the solvent phase.

15/21


FIGURE 15



Release profile of theophylline in simulated intestinal fluid from K6 (2% HPC) microspheres with a total polymer concentration of 7.5% in the solvent phase.

16/21


FIGURE 16



Release profile of theophylline in simulated intestinal fluid from K7 (5% HPC) microspheres with a total polymer concentration of 7.5% in the solvent phase.

17/21


FIGURE 17



Release profile of theophylline in simulated intestinal fluid from K5 (0.5% HPC) microspheres with a total polymer concentration of 7% in the solvent phase.

18/21


FIGURE 18



Release profile of theophylline in simulated intestinal fluid from K9 (2.5% HPC) microspheres with a total polymer concentration of 8% in the solvent phase.

19/21


FIGURE 19



Release profile of theophylline in simulated intestinal fluid from K18 (5% HPC) microspheres with a total polymer concentration of 9% in the solvent phase.

20/21


FIGURE 20



Release profiles from 180-250 μm size fraction microspheres for different microspheres preparations compared to that of CAB 381-20 alone.

21/21

FIGURE 21



Release profiles from 250-355µm size fraction microspheres for different microspheres preparations compared to that of CAB 381-20 alone.